



Anthephora ramosa on farm Onanis. (Photo: W. Giess)



Anthephora pubescens in the National Botanic Garden, Windhoek. (Photo: C. Mannheimer)



Spartina maritima at Lüderitz Bucht. (Photo: W. Giess)



Danthoniopsis sp. (Photo: W. Giess)



Rural people harvesting *Phragmites australis*, Caprivi. (Photo: C. Hines)



Elionurus muticus on farm Regenstein.
(Photo: W. Giess)



Eragrostis aristata at Orabes Quelle, Brandberg.
(Photo: H. Kolberg)



Entoplocamia aristulata, habit.
(Photo: C. Hines)



Melinis repens north of Otjikuare. (Photo: W. Giess)



Stipagrostis ciliata var. *capensis*, spreading hairs at the nodes.
(Photo: B. Loutit)



Aristida effusa at Otjitambi. (Photo: W. Giess)



Phragmites australis along the Okavango River. (Photo: W. Giess)



Stipagrostis plain in the Usakos area. (Photo: S. Schubert)



Pennisetum glaucum at Ondangwa. (Photo: W. Giess)



Stipagrostis giessii plain, Marienfluss, Kaokoveld. (Photo: W. Giess)



Odyssea paucinervis south of Hoanib River. (Photo: W. Giess)



Stipagrostis hirtigluma subsp. *pearsonii* on farm Otjitambi.
(Photo: W. Giess)



Melinis repens subsp. *grandiflora* at Otjitambi. (Photo: W. Giess)



Stipagrostis sabulicola on the way to Spencer Bay.
(Photo: C. Mannheimer)



Oryzidium barnardii at Ruacana Falls. (Photo: W. Giess)



Cenchrus ciliaris. (Photo: F. Bester)



Tiraphis ramisissima at Windhoek. (Photo: H. Kolberg)